



University of Groningen

## Measurement of the ratio of B-c(+) branching fractions to J/psi pi(+) and J/psi mu(+)nu(mu) final states

Aaij, R.; Adeva, B.; Adinolfi, M.; Affolder, A.; Ajaltouni, Z.; Akar, S.; Albrecht, J.; Alessio, F.; Alexander, M.; Ali, S.

*Published in:*  
Physical Review D

*DOI:*  
[10.1103/PhysRevD.90.032009](https://doi.org/10.1103/PhysRevD.90.032009)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2014

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Aaij, R., Adeva, B., Adinolfi, M., Affolder, A., Ajaltouni, Z., Akar, S., ... LHCb Collaboration (2014). Measurement of the ratio of B-c(+) branching fractions to J/psi pi(+) and J/psi mu(+)nu(mu) final states. Physical Review D, 90(3), [032009]. <https://doi.org/10.1103/PhysRevD.90.032009>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Measurement of the ratio of $B_c^+$ branching fractions to $J/\psi\pi^+$ and $J/\psi\mu^+\nu_\mu$ final states

R. Aaij *et al.*\*

(LHCb Collaboration)

(Received 8 July 2014; published 27 August 2014)

The first measurement that relates semileptonic and hadronic decay rates of the  $B_c^+$  meson is performed using proton-proton collision data corresponding to  $1.0 \text{ fb}^{-1}$  of integrated luminosity collected with the LHCb detector. The measured value of the ratio of branching fractions,  $\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi\mu^+\nu_\mu) = 0.0469 \pm 0.0028(\text{stat}) \pm 0.0046(\text{syst})$ , is at the lower end of available theoretical predictions.

DOI: 10.1103/PhysRevD.90.032009

PACS numbers: 12.39.Hg, 13.20.He, 13.25.Hw, 14.40.Nd

## I. INTRODUCTION

The  $B_c^+$  meson is the ground state of the  $\bar{b}c$  quark-pair system and is the only meson in which weak-interaction decays of both constituents compete with each other [1]. About 70% of the decay width is expected to be due to the  $c \rightarrow s$  transition, favored by the Cabibbo–Kobayashi–Maskawa quark-coupling hierarchy [2]. This decay process has recently been observed in the  $B_c^+ \rightarrow B_s^0\pi^+$  mode [3]. The complementary  $b \rightarrow c$  transition, which is predicted to account for 20% of the decay width, is more straightforward to observe experimentally, having a substantial probability to produce a  $J/\psi$  meson. Among such decays, semileptonic  $B_c^+ \rightarrow J/\psi\ell^+\nu_\ell$  ( $\ell = \mu, e$ ) and hadronic  $B_c^+ \rightarrow J/\psi\pi^+$  channels have played a special role in many measurements. The semileptonic decays were used in the discovery of the  $B_c^+$  meson [4], the measurements of its lifetime [4–7] and the measurement of the production cross section at the Tevatron [4]. The  $B_c^+ \rightarrow J/\psi\pi^+$  decays were used to measure its lifetime [8], mass [9–11], production cross section at the LHC [11] and as a reference for other hadronic branching fraction measurements [12–17]. However, there is no experimental determination of the relative size of semileptonic and hadronic decay rates. The goal of this work is a measurement of the ratio of branching fractions,

$$\mathcal{R} \equiv \frac{\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)}{\mathcal{B}(B_c^+ \rightarrow J/\psi\mu^+\nu_\mu)}, \quad (1)$$

and to test various theoretical models of  $B_c^+$  meson decays, for which predictions of  $\mathcal{R}$  vary over a wide range, 0.050–0.091 [18–25].

\*Full author list given at the end of the article.

Published by the American Physical Society under the terms of the Creative Commons Attribution 3.0 License. Further distribution of this work must maintain attribution to the author(s) and the published articles title, journal citation, and DOI.

## II. ANALYSIS OUTLINE

Final states containing a muon offer a distinctive experimental signature and can be triggered and reconstructed with high efficiency at LHCb. Therefore, this analysis relies on  $J/\psi$  decays to  $\mu^+\mu^-$ . Since the neutrino is not detected, both of the studied decay modes are reconstructed using a  $J/\psi$  candidate plus a charged track ( $\pi^+$ ), referred to as the *bachelor* track. The mass of  $J/\psi\pi^+$  signal candidates peaks at the  $B_c^+$  mass within the experimental resolution, allowing a straightforward signal-yield extraction in the presence of relatively small backgrounds under the signal peak. The main challenge in this analysis is the signal-yield extraction for the  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  decay mode, as the  $J/\psi\mu^+$  mass ( $m_{J/\psi\mu}$ ) distribution is broad due to the undetected neutrino. To suppress the dominant backgrounds, the analysis is restricted to the  $m_{J/\psi\mu} > 5.3 \text{ GeV}$  end point region and uses the mass-shape difference between the signal and the remaining background to extract the  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  signal yield [26]. In this mass region the neutrino has low energy; thus the  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  candidates are kinematically similar to the  $B_c^+ \rightarrow J/\psi\pi^+$  candidates. Therefore, many reconstruction uncertainties cancel in the ratio of their rates, allowing a precise measurement of  $\mathcal{R}(m_{J/\psi\mu} > 5.3 \text{ GeV})$ . This end point value is then extrapolated to the full phase space using theoretical predictions. Since the  $B_c^+$  and  $J/\psi$  are both  $1S$  heavy quarkonia states, the form factors involved in predicting the extrapolation factor and the shape of the mass distribution at the end point have only modest model dependence.

## III. DETECTOR AND DATA SAMPLE

The analysis is performed on a data sample of  $pp$  collisions at a center-of-mass energy of 7 TeV, collected during 2011 by the LHCb experiment and corresponding to an integrated luminosity of  $1.0 \text{ fb}^{-1}$ . The LHCb detector [27] is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , designed for the study of particles containing  $b$  or  $c$  quarks. The detector includes a

high-precision tracking system consisting of a silicon-strip vertex detector surrounding the  $pp$  interaction region [28], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [29] placed downstream of the magnet. The tracking system provides a measurement of momentum,  $p$ , with a relative uncertainty that varies from 0.4% at low momentum to 0.6% at 100 GeV. The minimum distance of a track to a primary vertex, the impact parameter (IP), is measured with a resolution of  $(15 + 29/p_T) \mu\text{m}$ , where  $p_T$  is the component of  $p$  transverse to the beam, in GeV. Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors [30]. Photon, electron and hadron candidates are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [31].

Simulated event samples are generated for the signal decays and the decay modes contributing to the background. In the simulation,  $pp$  collisions are generated using PYTHIA [32] with a specific LHCb configuration [33]. The production of  $B_c^+$  mesons, which is not adequately simulated in PYTHIA, is performed by the dedicated generator BCVEGPY [34]. Several dynamical models are used to simulate  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  decays. Decays of hadronic particles are described by EVTGEN [35], in which final-state radiation is generated using PHOTOS [36]. The interaction of the generated particles with the detector and its response are implemented using the GEANT4 toolkit [37] as described in Ref. [38].

#### IV. DATA SELECTION

This analysis relies on  $J/\psi t^+$  candidates satisfying the trigger [39], which consists of a hardware stage, based on information from the muon system, followed by a two-level software stage, which applies a full event reconstruction. At the hardware stage, a muon with  $p_T > 1.5$  GeV, or a pair of muons with  $\sqrt{p_{T1} p_{T2}} > 1.3$  GeV, is required. The subsequent lower-level software triggers require a charged-particle track with  $p_T > 1.7$  GeV ( $p_T > 1.0$  GeV if identified as muon) and with an IP relative to any primary  $pp$ -interaction vertex (PV) larger than 100  $\mu\text{m}$ . A dimuon trigger, which requires a large dimuon mass,  $m_{\mu^+\mu^-} > 2.7$  GeV, and each muon to have  $p_T > 0.5$  GeV, complements the single track triggers. The final software trigger stage requires either a  $J/\psi \rightarrow \mu^+\mu^-$  candidate with a  $J/\psi$  decay vertex separation from the nearest PV of at least three standard deviations, or that a two- or three-track combination, which includes a muon, is identified as a secondary vertex using a multivariate selection [39].

In the offline analysis,  $J/\psi \rightarrow \mu^+\mu^-$  candidates are selected with the following criteria:  $p_T(\mu) > 0.9$  GeV,

$p_T(J/\psi) > 1.5$  GeV,  $\chi^2$  per degree of freedom (ndf) for the two muons to form a common vertex  $\chi_{\text{vtx}}^2(\mu^+\mu^-)/\text{ndf} < 9$ , and a mass consistent with the  $J/\psi$  meson. The separation of the  $J/\psi$  decay vertex from the nearest PV must be at least five standard deviations. The bachelor track, and at least one of the muons from the decay of the  $J/\psi$  meson, must not point to any PV, through the requirement  $\chi_{\text{IP}}^2 > 9$ . The quantity  $\chi_{\text{IP}}^2$  is defined as the difference between the  $\chi^2$  of the PV fitted with and without the considered particle. The bachelor track must not be collinear within  $0.8^\circ$  with either of the muons from the  $J/\psi$  meson decay and must satisfy  $p_T > 0.5$  GeV ( $> 1.0$  GeV for  $\pi^+$ ). A loose kaon veto is applied to the pion candidates,  $\ln[\mathcal{L}(K)/\mathcal{L}(\pi)] < 5$ , where  $\mathcal{L}$  is the particle identification likelihood [40]. The  $J/\psi$  candidates are combined with the bachelor tracks in a kinematic fit to form  $B_c^+$  candidates with the known  $J/\psi$  mass and the  $B_c^+$  vertex used as constraints. The  $B_c^+$  candidate must satisfy  $\chi_{\text{vtx}}^2(J/\psi t^+)/\text{ndf} < 9$  and have a pseudoproper decay time greater than 0.25 ps. The pseudoproper decay time is determined as  $L \cdot m_{J/\psi t^+}/|\vec{p}_{J/\psi t^+}|$ , where  $L$  is the projection of the distance between the  $B_c^+$  production and decay vertices onto the direction of the  $J/\psi t^+$  momentum  $\vec{p}_{J/\psi t^+}$  and  $m_{J/\psi t^+}$  is the  $J/\psi t^+$  mass.

Four discriminating variables ( $x_i$ ) are used in a likelihood ratio to improve the background suppression. Three of the variables are common between the two channels:  $\chi_{\text{vtx}}^2(J/\psi t^+)/\text{ndf}$ ,  $\chi_{\text{IP}}^2(B_c^+)$ , and the cosine of the angle between the  $J/\psi$  meson and the bachelor track transverse momenta. The latter quantity peaks at positive values for the signal as the  $B_c^+$  meson has a high transverse momentum. Background events in which particles are combined from two different  $B$  decays usually peak at negative values, while those due to random combinations of particles are more uniformly distributed. The  $\chi_{\text{IP}}^2(B_c^+)$  variable is small for  $B_c^+ \rightarrow J/\psi \pi^+$  decays since the  $B_c^+$  momentum points back to the PV. For  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  candidates, the pointing is only approximate since the neutrino is not reconstructed. However,  $\chi_{\text{IP}}^2(B_c^+)$  is often smaller than for the background events because the neutrino has low momentum. The fourth variable for the  $J/\psi \pi^+$  mode is  $\chi_{\text{IP}}^2(t^+)$ , while for the  $J/\psi \mu^+ \nu_\mu$  mode it is the pseudoproper decay time, as  $\chi_{\text{IP}}^2(t^+)$  is found to be ineffective for this channel. The four one-dimensional signal probability density functions (PDFs),  $\mathcal{P}_{\text{sig}}(x_i)$ , are obtained from a simulated sample of signal events. The background PDFs,  $\mathcal{P}_{\text{bkg}}(x_i)$ , are obtained from the data in the  $B_c^+ \rightarrow J/\psi \pi^+$  mass sidebands (5.35–5.80 and 6.80–8.50 GeV) and from the simulation of inclusive backgrounds from  $B_{u,d,s} \rightarrow J/\psi X$  decays ( $X$  denotes one or more particles) for the  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  candidates. The requirement  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L}) = -2 \sum_{i=1}^4 \ln[\mathcal{P}_{\text{sig}}(x_i)/\mathcal{P}_{\text{bkg}}(x_i)] < 1.0$  ( $< 0.0$ ) preserves about 93% (87%) of signal events for  $B_c^+ \rightarrow J/\psi \pi^+$  ( $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  with  $m_{J/\psi \mu} > 5.3$  GeV) and efficiently suppresses the backgrounds. These requirements

minimize the expected average statistical uncertainty on the signal yields, given the observed background levels in each channel.

### V. EXTRACTION OF THE $B_c^+ \rightarrow J/\psi\pi^+$ SIGNAL

An extended maximum likelihood fit to the unbinned distribution of observed  $m_{J/\psi\pi}$  values yields  $N_{J/\psi\pi} = 839 \pm 40$   $B_c^+ \rightarrow J/\psi\pi^+$  signal events and is shown in Fig. 1.

The signal is represented in the fit by a double-sided Crystal Ball function [41]. The peak position, the Gaussian mass resolution and the peak amplitude are free parameters in the fit, while the parameters describing small non-Gaussian tails are fixed by a fit to the simulated signal distribution. Using a Gaussian function to model the signal results in a 2.3% relative change in  $\mathcal{R}$  value, and this is assigned as the systematic uncertainty. The background is smoothly distributed and modeled by an exponential function. Varying the background parametrization and the fit range results in up to a 0.6% relative change in  $\mathcal{R}$ . A small background from  $B_c^+ \rightarrow J/\psi K^+$  decays, peaking 37 MeV below the signal peak, is also included in the fit with all shape parameters fixed from the simulation. Its normalization is constrained to be 1% of the fitted signal amplitude, as predicted by the measured ratio of the branching fractions [15] scaled by an efficiency ratio of 15% obtained from the simulation. The relative systematic uncertainty on  $\mathcal{R}$  related to this fit component is 0.1%.

### VI. EXTRACTION OF THE $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$ SIGNAL

To measure the  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  rate, feed down from other  $B_c^+ \rightarrow f$ ,  $f \rightarrow J/\psi\mu^+\nu_\mu X$  decays must be accounted for. Decays to excited charmonium states [ $f = \psi_f\mu^+\nu_\mu$ ,




FIG. 1 (color online). Invariant-mass distribution of  $B_c^+ \rightarrow J/\psi\pi^+$  candidates (black data points). The maximum likelihood fit of the  $B_c^+$  signal is superimposed (blue solid line). Individual fit components are also shown: (dashed blue line) the signal, (red long-dashed line) the background and (green dotted line)  $B_c^+ \rightarrow J/\psi K^+$  feed down.

with  $\psi_f = \chi_{cJ}$  or  $\psi(2S)$ ] and states containing  $\tau$  leptons ( $f = J/\psi\tau^+\nu_\tau$ ) are the dominant contributions. Since the rates for such decays have not been measured, we rely on theoretical predictions for

$$R_f \equiv \frac{\mathcal{B}(B_c^+ \rightarrow f)}{\mathcal{B}(B_c^+ \rightarrow J/\psi\mu^+\nu_\mu)}. \quad (2)$$

Although the spread in  $R_f$  predictions is large (see below), the related systematic uncertainty is minimized by restricting the analysis to the high  $J/\psi\mu^+$  mass region. Unreconstructed decay products in the  $\psi_f \rightarrow J/\psi X$  transitions ( $X = \gamma, \pi\pi, \pi^0, \eta, \gamma\gamma$ ) or  $\tau^+ \rightarrow \mu^+\nu_\mu\bar{\nu}_\tau$  decays carry energy away, lowering the  $J/\psi\mu^+$  mass relative to that from direct  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  decays, as illustrated in Fig. 2. The selection requirement in  $m_{J/\psi\mu}$  is chosen to eliminate the backgrounds from  $B_{u,d,s}$  decays to  $J/\psi$  mesons associated with hadrons, with one of the hadrons misidentified as a muon. These backgrounds are large because the  $B_{u,d,s}$  production rates are orders of magnitude higher than for  $B_c^+$ . Since many exclusive decay modes with various hadron multiplicities and unknown branching ratios contribute, the  $m_{J/\psi\mu}$  shape of such backgrounds is difficult to predict. The 5.3 GeV lower limit on  $m_{J/\psi\mu}$  is above the kinematic limit for  $B_u^+ \rightarrow J/\psi h^+$  decays, with  $h^+$  denoting a charged kaon or pion, as illustrated in Fig. 2. The  $B_{u,d,s}$  backgrounds in the selected region are much smaller, and are from  $B_{u,d,s} \rightarrow J/\psi X$  decays paired with a bachelor  $\mu^+$  originating from a semileptonic decay of the companion  $b$  quark in the produced  $b\bar{b}$  pair. Simulation of  $b$ -baryon decays to final states involving a  $J/\psi$  meson shows that they also contribute via this mechanism. The shape of such




FIG. 2 (color online). Distribution of  $m_{J/\psi\mu}$  for  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  candidates selected in simulated event samples of (blue filled points) the signal, (green filled points) the  $B_c^+$  feed down and (red filled squares) the  $B_{u,d,s}$  backgrounds. Relative normalization is derived from the fit to the data described later in the text. The part of the spectrum included in the fit is indicated with a vertical dashed black line. The  $B_c^+$  feed down distribution is also shown after magnifying its normalization by a factor of 10 (green dashed histogram).



combinatorial backgrounds is less sensitive to the details of the composition of  $b$ -hadron decay modes, and thus is easier to predict. Since the combinatorial backgrounds are dominated by genuine muons, the analysis is not sensitive to the estimation of muon misidentification rates and associated systematic uncertainties.

The  $m_{J/\psi\mu}$  signal shape is dominated by the end point kinematics, whereas the combinatorial background is smooth and extends beyond the kinematic limit for the  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  decays. The signal yield is determined by a fit to the  $m_{J/\psi\mu}$  distribution. The feed down background is small as discussed in detail below. Its shape is constrained by simulation, while its normalization is related to the signal yield via theoretical predictions. The unbinned maximum likelihood fit is performed simultaneously to the  $m_{J/\psi\mu}$  distribution in data and the signal and background distributions from simulation, in the range of 5.3 to 8.0 GeV, and gives  $N_{J/\psi\mu} = 3537 \pm 125$  signal events. The  $m_{J/\psi\mu}$  distributions and the fit results are displayed in Fig. 3. The fit is described in detail below.

The total PDF used in the fit is the sum of the signal PDF ( $\mathcal{P}_{\text{sig}}$ ), the feed down background PDF ( $\mathcal{P}_{\text{fd}}$ ) and the combinatorial background PDF ( $\mathcal{P}_{\text{bkg}}$ ),

$$\mathcal{P}(m_{J/\psi\mu}) \propto N_{J/\psi\mu}(\mathcal{P}_{\text{sig}}(m_{J/\psi\mu}) + \alpha\mathcal{P}_{\text{fd}}(m_{J/\psi\mu})) + N_{\text{bkg}}\mathcal{P}_{\text{bkg}}(m_{J/\psi\mu}), \quad (3)$$

where  $\alpha$  is the feed-down-to-signal yield ratio and  $N_{\text{bkg}}$  is the combinatorial background yield. The signal shape is dominated by the end point kinematics; thus it is modeled as

$$\mathcal{P}_{\text{sig}}(m_{J/\psi\mu}) \propto \text{PS}(m_{J/\psi\mu})(1 + s_1\bar{m}_{J/\psi\mu}), \quad (4)$$

where  $\bar{m}_{J/\psi\mu} = m_{J/\psi\mu} - 5.3$  GeV and  $\text{PS}(m_{J/\psi\mu})$  corresponds to the uniform distribution in the  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  three-body phase-space,

$$\begin{aligned} \text{PS}(m_{J/\psi\mu}) &= \frac{M_{B_c}^2 - m_{J/\psi\mu}^2}{m_{J/\psi\mu}} \\ &\times \sqrt{m_{J/\psi\mu}^2 - (M_{J/\psi} + M_\mu)^2} \\ &\times \sqrt{m_{J/\psi\mu}^2 - (M_{J/\psi} - M_\mu)^2}, \end{aligned} \quad (5)$$

with the  $J/\psi$  and  $\mu$  masses ( $M_{J/\psi}$  and  $M_\mu$ ) set to their known values [42] and  $M_{B_c}$  set to an effective value, which




FIG. 3 (color online). Invariant-mass distribution of  $J/\psi\mu^+$  pairs from  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  candidates (black data points) for (top left) the data, (bottom left)  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  signal simulation, (top right)  $B_{u,d,s} \rightarrow J/\psi X$  background simulation and (bottom right)  $B_c^+$  feed down simulation. The unbinned maximum likelihood fit of the  $B_c^+$  signal is superimposed (blue solid line). Individual fit components are also shown: (blue short-dashed line) the signal, (red long-dashed line) the background and (green dotted line)  $B_c^+$  feed down.

is slightly higher than the  $B_c^+$  mass to account for detector resolution effects. Setting  $M_{B_c}$  to the known  $B_c^+$  mass [42] changes the signal yield by a negligible amount. Deviations from the uniform distribution are allowed by the linear term, with the  $s_1$  coefficient determined by the simultaneous fit to the simulated signal distribution and the data. The simulation based on the Kiselev *et al.* QCD sum rules model [22] is used in the default fit. The models of Ebert *et al.* [23], based on a relativistic quasipotential Schrödinger approach, and ISGW2 [43], based on a non-relativistic constituent quark model with relativistic corrections, alter the determined signal yield by +0.2% and -0.4%, respectively. Relying on the data themselves to determine the signal shape changes the signal yield by +0.7%. The latter value is taken as a systematic error.

The feed down includes contributions from the following  $B_c^+$  decay modes  $f = \psi(2S)\mu^+\nu_\mu$ ,  $\chi_{cJ}\mu^+\nu_\mu$  and  $J/\psi\tau^+\nu_\tau$ . Feed down from  $B_c^+ \rightarrow B_{d,s}\mu^+\nu_\mu$  and  $B_c^+ \rightarrow J/\psi$  plus hadrons is also investigated and found negligible. Their individual proportions with respect to the signal yield are determined as

$$\alpha_f = R_f \mathcal{B}_{\text{casc } f} R_{ef}, \quad (6)$$

and then added,  $\alpha = \sum_f \alpha_f$ , where  $\mathcal{B}_{\text{casc } f}$  is the sum of the measured branching fractions [42] for the  $\psi_f$  state to decay to a  $J/\psi$  meson by emission of unreconstructed photons or light hadrons, and  $R_{ef}$  is the ratio of the feed down and the signal reconstruction efficiencies [44]. This quantity is small because of the  $m_{J/\psi\mu} > 5.3$  GeV requirement. For  $\chi_{cJ}$  states the sum extends over the three  $J$  values,  $R_f \mathcal{B}_{\text{casc } f} = \sum_{J=0,1,2} R_{fJ} \mathcal{B}_{\text{casc } fJ}$ . The values of the parameters affecting the estimate of the feed down fraction are summarized in Table I. Theoretical predictions for  $R_f$  for the  $B_c^+ \rightarrow \psi(2S)\mu^+\nu_\mu$  feed down mode vary over a wide range, 0.009–0.185 [18,21–23,25,43,45]. An average of the highest and the lowest prediction is taken for the nominal estimate, and half of the difference is taken for the systematic error. The theoretical uncertainties in the  $R_f \mathcal{B}_{\text{casc } f}$  values for the dominant  $B_c^+ \rightarrow \chi_{cJ}\mu^+\nu_\mu$  feed down mode are smaller, 0.032–0.038 [24,46,47]. The spread is also limited for theoretical predictions of  $R_f$  for the  $B_c^+ \rightarrow J/\psi\tau^+\nu_\tau$  decay, 0.237–0.283 [19,22,24,47]. The simulated distributions for the individual feed down modes are mixed according to the proportions resulting from the  $R_f \mathcal{B}_{\text{casc } f}$  values and then parametrized as

$$\mathcal{P}_{\text{fd}}(m_{J/\psi\mu}) \propto \text{PS}(m_{J/\psi\mu})(1 + f_1 \bar{m}_{J/\psi\mu} + f_2 \bar{m}_{J/\psi\mu}^2), \quad (7)$$

where  $f_1$  and  $f_2$  are parameters determined by the fit. The effect of the unreconstructed decay products  $X$  is to lower the effective  $M_{B_c}$  value in Eq. (5). Varying the feed down fraction within its uncertainty changes the signal yield by up to 0.6%.

The combinatorial  $B_{u,d,s}$  background is parametrized with an exponential function. The tail of the  $B_u^+ \rightarrow J/\psi h^+$  distribution, with the light hadron misidentified as a muon, may enter the signal region because of detector resolution. We parametrize it with a Gaussian function,  $G(m_{J/\psi\mu})$ , with a mean value and width fixed to the results of the fit to the simulated  $B_u^+ \rightarrow J/\psi h^+$  distribution. The exponential and  $G(m_{J/\psi\mu})$  functions together define  $\mathcal{P}_{\text{bkg}}(m_{J/\psi\mu})$ ,

$$\mathcal{P}_{\text{bkg}}(m_{J/\psi\mu}) \propto c N_e e^{b_1 \bar{m}_{J/\psi\mu} + b_2 \bar{m}_{J/\psi\mu}^2} + (1 - c)G(m_{J/\psi\mu}),$$

where  $N_e$  normalizes the exponential function to 1. The combinatorial background fraction  $c$  and the polynomial coefficients  $b_1$  and  $b_2$  are free parameters in the simultaneous fit to the simulated  $B_{u,d,s} \rightarrow J/\psi X$  distribution and to the distribution in the data. To avoid relying on simulation for the absolute values of the muon misidentification rates,  $c$  is allowed to vary independently in the fit to the simulated and the observed distributions. A systematic uncertainty of 1.8% is assigned to this background parametrization based on fit results in which either the Gaussian term is neglected or the exponential function is replaced by a sum of two exponential functions.

Varying the upper limit of the mass range used in the fit from 8.0 down to 6.75 GeV results in a signal-yield change of up to 1.5%. Varying the corresponding lower limit from its default value of 5.3 to 5.1 GeV, thus including the peak of the  $B_u^+ \rightarrow J/\psi h^+$  component (see Fig. 2), or to 5.5 GeV, thus avoiding the tail of that component, results in a relative change in the  $\mathcal{R}$  value of up to 1.6%.

The default method of the  $B_c^+ \rightarrow J/\psi\mu^+\nu_\mu$  signal-yield determination relies on simulation to predict the signal and background shapes in the  $m_{J/\psi\mu}$  distribution. An alternative approach relies on simulation to predict the signal and background shapes of the  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  distribution. Correlations between  $m_{J/\psi\mu}$  and  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  variables are small. The requirement on the  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  value is removed. The  $m_{J/\psi\mu}$  range is restricted to 5.3–6.1 GeV to exclude the backgrounds above the  $B_c^+$  kinematic limit.

TABLE I. Values of the parameters affecting the estimate of the feed down fraction in the fit to the  $m_{J/\psi\mu}$  distribution. For  $B_c^+ \rightarrow \chi_{cJ}\mu^+\nu_\mu$ ,  $\sum_{J=0,1,2} R_{fJ} \mathcal{B}_{\text{casc } fJ}$  is listed.

Feed down mode	$R_f$	$\mathcal{B}_{\text{casc } f}$	$R_{ef}$	$\alpha_f$
$B_c^+ \rightarrow \psi(2S)\mu^+\nu_\mu$	0.009–0.185	$0.598 \pm 0.006$	$0.118 \pm 0.004$	$0.0069 \pm 0.0062$
$B_c^+ \rightarrow \chi_{cJ}\mu^+\nu_\mu$		0.032–0.038	$0.364 \pm 0.009$	$0.0127 \pm 0.0011$
$B_c^+ \rightarrow J/\psi\tau^+\nu_\tau$	0.237–0.283	$0.1741 \pm 0.0004$	$0.014 \pm 0.001$	$0.0006 \pm 0.0001$
Total $\alpha$				$0.0202 \pm 0.0063$

The signal and combinatorial background yields are determined by a fit to the  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  distribution in the data. The  $B_c^+$  feed down simulation predicts a similar  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  shape as for the  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  signal. Therefore, this contribution is not represented explicitly in the fit to the  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  distribution, but is subtracted from the fitted signal yield according to the feed down fraction  $\alpha$ . Taking into account the differences in signal efficiency, the  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  signal yield is consistent with that resulting from the  $m_{J/\psi \mu}$  fit method within 0.5%, which is included as an additional systematic uncertainty due to the  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  requirement in the nominal approach.

## VII. RESULTS

The ratio of the reconstruction efficiencies between the two  $B_c^+$  signal modes, as determined from simulation, is  $\epsilon(B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu)/\epsilon(B_c^+ \rightarrow J/\psi \pi^+) = 1.14 \pm 0.01$  (statistical error) for  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  events generated in the end point region. Using different  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  form factor models changes this efficiency ratio by up to 1.3%. Efficiencies of the pion and muon particle identification (PID) requirements have systematic uncertainties of 0.8% and 1.9%, respectively. The efficiency-ratio systematic uncertainties from the  $B_c^+$  lifetime assumed in the simulation is 0.2% due to the cancelations between the two decay modes. The fraction of multiple signal candidates per event is 0.1% for  $B_c^+ \rightarrow J/\psi \pi^+$  and 1.9% for  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  decays. To check for possible biases due to the neglected correlations between multiple candidates, one candidate is randomly chosen, which changes the  $\mathcal{R}$  result by 0.4%. The systematic uncertainty associated with the limited knowledge of the efficiency of the  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  requirement for  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  decays is included using the results of the  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  fit. To study the corresponding uncertainty for  $B_c^+ \rightarrow J/\psi \pi^+$  decays, the  $\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$  requirement is varied, resulting in a 2% variation. The systematic uncertainty associated with the trigger simulation is 3.4%, as estimated by modifying the trigger requirements. The systematic errors are summarized in Table II. The total relative systematic uncertainty on  $\mathcal{R}(m_{J/\psi \mu} > 5.3 \text{ GeV})$  is 6%.

The result for the ratio of the branching fractions restricted to decays with  $m_{J/\psi \mu} > 5.3 \text{ GeV}$  is

$$\mathcal{R}(m_{J/\psi \mu} > 5.3 \text{ GeV}) = 0.271 \pm 0.016 \pm 0.016, \quad (8)$$

where the first uncertainty is statistical and the second is systematic. This ratio is extrapolated to the full phase space using the predictions of the phenomenological models [22,23,25,43,48,49]. No correlation pattern is observed between the predicted values of  $\mathcal{R}$  and of the extrapolation factor; thus equal weight is given to all models. The model of Kiselev *et al.* [22] predicts the fraction of the  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$  rate with  $m_{J/\psi \mu}$  above 5.3 GeV to be 0.173,

TABLE II. Summary of systematic uncertainties. The total systematic errors are obtained by adding in quadrature the individual contributions.

Contribution	Relative error
$m_{J/\psi \pi}$ signal shape	2.3%
$m_{J/\psi \pi}$ background shape	0.2%
$B_c^+ \rightarrow J/\psi K^+$ component	0.1%
$m_{J/\psi \mu}$ signal shape	0.7%
$m_{J/\psi \mu}$ background shape	1.8%
$B_c^+$ feed down	0.6%
Lower $m_{J/\psi \mu}$ fit range limit	1.6%
Upper $m_{J/\psi \mu}$ fit range limit	1.5%
$B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$ model dependence of efficiency	1.3%
Pion PID	0.8%
Muon PID	1.9%
Lifetime	0.2%
Multiple candidates	0.4%
$\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$ requirement for $B_c^+ \rightarrow J/\psi \pi^+$	2.0%
$\Delta_{\text{sig/bkg}}(-2 \ln \mathcal{L})$ requirement for $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu$	0.5%
Trigger simulation	3.4%
Total within selected $m_{J/\psi \mu}$ range	6.0%
$m_{J/\psi \mu}$ extrapolation	7.9%
Total	9.9%

which is close to the average over all models. The largest deviation from this prediction is 7.9%, which is taken as an estimate of the extrapolation systematic error. This increases the systematic uncertainty on  $\mathcal{R}$ , when extrapolated to the full mass range, to 9.9% yielding




FIG. 4 (color online). The measured value of  $\mathcal{R}$  (horizontal solid line) and its  $\pm 1\sigma$  uncertainty band (dashed lines) compared to the predictions (diamonds). A nonrelativistic reduction of the Bethe–Salpeter equation is used in the predictions of Chang *et al.* [18], El-Hady *et al.* [20] and Colangelo *et al.* [21], while the latter also utilizes heavy quark symmetry. A light-front constituent quark model is used by Anisimov *et al.* [19] and Ke *et al.* [25]. QCD sum rules are used by Kiselev *et al.* [22], a relativistic quasipotential Schrödinger model is used by Ebert *et al.* [23], and a relativistic constituent quark model is used by Ivanov *et al.* [24].

$$\mathcal{R} = 0.0469 \pm 0.0028 \pm 0.0046. \quad (9)$$

A comparison between the measured and the predicted values of  $\mathcal{R}$  is shown in Fig. 4. The measured value is slightly below the lowest predicted value. The predictions by the relativistic quasipotential Schrödinger model of Ebert *et al.* [23] and the model of El-Hady *et al.*, based on a nonrelativistic reduction of the Bethe–Salpeter equation [20], are in good agreement with the experimental value. The model of Ke *et al.* [25], based on the modified harmonic oscillator wave function in light-front quark model, is also consistent with the data. The other models [18,19,21,22,24] significantly overestimate  $\mathcal{R}$ .

### VIII. SUMMARY

The ratio of hadronic and semileptonic decay branching fractions of the  $B_c^+$  meson is measured for the first time. Within the observed mass range,  $m_{J/\psi\mu} > 5.3$  GeV, the measured value of  $\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi\mu^+\nu_\mu)$  is found to be  $0.271 \pm 0.016(\text{stat}) \pm 0.016(\text{syst})$ . Extrapolating to the full mass range, we obtain a value of  $\mathcal{B}(B_c^+ \rightarrow J/\psi\pi^+)/\mathcal{B}(B_c^+ \rightarrow J/\psi\mu^+\nu_\mu) = 0.0469 \pm 0.0028(\text{stat}) \pm 0.0046(\text{syst})$ , which is in good agreement with the theoretical predictions by Ebert *et al.* [23] and El-Hady *et al.* [20], and consistent with the prediction by Ke *et al.* [25]. All other currently available models [18,19,21,22,24] overestimate this ratio.

### ACKNOWLEDGMENTS

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG, HGF and MPG (Germany); SFI (Ireland); INFN (Italy); FOM and NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FANO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); and NSF (USA). The Tier1 computing centers are supported by IN2P3 (France), KIT and BMBF (Germany), INFN (Italy), NWO and SURF (Netherlands), PIC (Spain) and GridPP (United Kingdom). We are indebted to the communities behind the multiple open source software packages on which we depend. We are also thankful for the computing resources and the access to software research and development tools provided by Yandex LLC (Russia). Individual groups or members have received support from EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union); Conseil général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France); RFBR (Russia); XuntaGal and GENCAT (Spain); and Royal Society and Royal Commission for the Exhibition of 1851 (United Kingdom).

- 
- [1] Charge-conjugate states are implied in this article.
  - [2] I. P. Gouz, V. V. Kiselev, A. K. Likhoded, V. I. Romanovsky, and O. P. Yushchenko, *Phys. At. Nucl.* **67**, 1559 (2004).
  - [3] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **111**, 181801 (2013).
  - [4] F. Abe *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **81**, 2432 (1998).
  - [5] A. Abulencia *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **97**, 012002 (2006).
  - [6] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **102**, 092001 (2009).
  - [7] R. Aaij *et al.* (LHCb Collaboration), *Eur. Phys. J. C* **74**, 2839 (2014).
  - [8] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **87**, 011101 (2013).
  - [9] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **100**, 182002 (2008).
  - [10] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **101**, 012001 (2008).
  - [11] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **109**, 232001 (2012).
  - [12] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. Lett.* **108**, 251802 (2012).
  - [13] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. D* **87**, 071103 (2013).
  - [14] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. D* **87**, 112012 (2013).
  - [15] R. Aaij *et al.* (LHCb Collaboration), *J. High Energy Phys.* **09** (2013) 075.
  - [16] R. Aaij *et al.* (LHCb Collaboration), *J. High Energy Phys.* **11** (2013) 094.
  - [17] R. Aaij *et al.* (LHCb Collaboration), *J. High Energy Phys.* **05** (2014) 148.
  - [18] C.-H. Chang and Y.-Q. Chen, *Phys. Rev. D* **49**, 3399 (1994).
  - [19] A. Y. Anisimov, I. M. Narodetskii, C. Semay, and B. Silvestre-Blac, *Phys. Lett. B* **452**, 129 (1999); A. Y. Anisimov, P. Y. Kulikov, I. M. Narodetskii, and K. A. Ter-Martirosyan, *Phys. At. Nucl.* **62**, 1739 (1999).
  - [20] A. A. El-Hady, J. H. Muñoz, and J. P. Vary, *Phys. Rev. D* **62**, 014019 (2000).
  - [21] P. Colangelo and F. De Fazio, *Phys. Rev. D* **61**, 034012 (2000).
  - [22] V. V. Kiselev, *arXiv:hep-ph/0211021*.
  - [23] D. Ebert, R. N. Faustov, and V. O. Galkin, *Phys. Rev. D* **68**, 094020 (2003).
  - [24] M. A. Ivanov, J. G. Körner, and P. Santorelli, *Phys. Rev. D* **73**, 054024 (2006).



- [25] H.-W. Ke, T. Liu, and X.-Q. Li, *Phys. Rev. D* **89**, 017501 (2014).  
 [26] Units in which  $c = 1$  are used.  
 [27] A. A. Alves, Jr. *et al.* (LHCb Collaboration), *J. Instrum.* **3**, S08005 (2008).  
 [28] R. Aaij *et al.*, [arXiv:1405.7808](#).  
 [29] R. Arink *et al.*, *J. Instrum.* **9**, P01002 (2014).  
 [30] M. Adinolfi *et al.*, *Eur. Phys. J. C* **73**, 2431 (2013).  
 [31] A. A. Alves, Jr. *et al.*, *J. Instrum.* **8**, P02022 (2013).  
 [32] T. Sjöstrand, S. Mrenna, and P. Skands, *J. High Energy Phys.* **05** (2006) 026.  
 [33] I. Belyaev *et al.*, Nuclear Science Symposium Conference Record (NSS/MIC) **IEEE**, 1155 (2010).  
 [34] C.-H. Chang, C. Driouichi, P. Eerola, and X. G. Wu, *Comput. Phys. Commun.* **159**, 192 (2004).  
 [35] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).  
 [36] P. Golonka and Z. Was, *Eur. Phys. J. C* **45**, 97 (2006).  
 [37] J. Allison *et al.* (GEANT4 Collaboration), *IEEE Trans. Nucl. Sci.* **53**, 270 (2006); S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).  
 [38] M. Clemencic, G. Corti, S. Easo, C. R. Jones, S. Miglioranza, M. Pappagallo, and P. Robbe, *J. Phys. Conf. Ser.* **331**, 032023 (2011).  
 [39] R. Aaij *et al.*, *J. Instrum.* **8**, P04022 (2013).  
 [40] M. Adinolfi *et al.*, *Eur. Phys. J. C* **73**, 2431 (2013).  
 [41] T. Skwarnicki, Ph.D. thesis, Institute of Nuclear Physics, Krakow, 1986 [Report No. DESY-F31-86-02].  
 [42] J. Beringer *et al.* (Particle Data Group), *Phys. Rev. D* **86**, 010001 (2012), and 2013 partial update for the 2014 edition.  
 [43] D. Scora and N. Isgur, *Phys. Rev. D* **52**, 2783 (1995).  
 [44] For  $B_c^+ \rightarrow J/\psi \tau^+ \nu_\tau$ ,  $\mathcal{B}_{\text{case } f} = \mathcal{B}(\tau^+ \rightarrow \mu^+ \nu_\mu \bar{\nu}_\tau)$ .  
 [45] J.-F. Liu and K.-T. Chao, *Phys. Rev. D* **56**, 4133 (1997).  
 [46] C.-H. Chang, Y.-Q. Chen, G.-L. Wang, and H.-S. Zong, *Phys. Rev. D* **65**, 014017 (2001).  
 [47] E. Hernandez, J. Nieves, and J. M. Verde-Velasco, *Phys. Rev. D* **74**, 074008 (2006).  
 [48] W. Wang, Y.-L. Shen, and C.-D. Lü, *Phys. Rev. D* **79**, 054012 (2009).  
 [49] W.-F. Wang, Y.-Y. Fan, and Z.-J. Xiao, [arXiv:1212.5903](#).

---

R. Aaij,<sup>41</sup> B. Adeva,<sup>37</sup> M. Adinolfi,<sup>46</sup> A. Affolder,<sup>52</sup> Z. Ajaltouni,<sup>5</sup> S. Akar,<sup>6</sup> J. Albrecht,<sup>9</sup> F. Alessio,<sup>38</sup> M. Alexander,<sup>51</sup> S. Ali,<sup>41</sup> G. Alkhazov,<sup>30</sup> P. Alvarez Cartelle,<sup>37</sup> A. A. Alves Jr.,<sup>25,38</sup> S. Amato,<sup>2</sup> S. Amerio,<sup>22</sup> Y. Amhis,<sup>7</sup> L. An,<sup>3</sup> L. Anderlini,<sup>17,a</sup> J. Anderson,<sup>40</sup> R. Andreassen,<sup>57</sup> M. Andreotti,<sup>16,b</sup> J. E. Andrews,<sup>58</sup> R. B. Appleby,<sup>54</sup> O. Aquines Gutierrez,<sup>10</sup> F. Archilli,<sup>38</sup> A. Artamonov,<sup>35</sup> M. Artuso,<sup>59</sup> E. Aslanides,<sup>6</sup> G. Auriemma,<sup>25,c</sup> M. Baalouch,<sup>5</sup> S. Bachmann,<sup>11</sup> J. J. Back,<sup>48</sup> A. Badalov,<sup>36</sup> V. Balagura,<sup>31</sup> W. Baldini,<sup>16</sup> R. J. Barlow,<sup>54</sup> C. Barschel,<sup>38</sup> S. Barsuk,<sup>7</sup> W. Barter,<sup>47</sup> V. Batozskaya,<sup>28</sup> V. Battista,<sup>39</sup> A. Bay,<sup>39</sup> L. Beaucourt,<sup>4</sup> J. Beddow,<sup>51</sup> F. Bedeschi,<sup>23</sup> I. Bediaga,<sup>1</sup> S. Belogurov,<sup>31</sup> K. Belous,<sup>35</sup> I. Belyaev,<sup>31</sup> E. Ben-Haim,<sup>8</sup> G. Bencivenni,<sup>18</sup> S. Benson,<sup>38</sup> J. Benton,<sup>46</sup> A. Berezhnoy,<sup>32</sup> R. Bernet,<sup>40</sup> M.-O. Bettler,<sup>47</sup> M. van Beuzekom,<sup>41</sup> A. Bien,<sup>11</sup> S. Bifani,<sup>45</sup> T. Bird,<sup>54</sup> A. Bizzeti,<sup>17,d</sup> P. M. Bjørnstad,<sup>54</sup> T. Blake,<sup>48</sup> F. Blanc,<sup>39</sup> J. Blouw,<sup>10</sup> S. Blusk,<sup>59</sup> V. Bocci,<sup>25</sup> A. Bondar,<sup>34</sup> N. Bondar,<sup>30,38</sup> W. Bonivento,<sup>15,38</sup> S. Borghi,<sup>54</sup> A. Borgia,<sup>59</sup> M. Borsato,<sup>7</sup> T. J. V. Bowcock,<sup>52</sup> E. Bowen,<sup>40</sup> C. Bozzi,<sup>16</sup> T. Brambach,<sup>9</sup> J. van den Brand,<sup>42</sup> J. Bressieux,<sup>39</sup> D. Brett,<sup>54</sup> M. Britsch,<sup>10</sup> T. Britton,<sup>59</sup> J. Brodzicka,<sup>54</sup> N. H. Brook,<sup>46</sup> H. Brown,<sup>52</sup> A. Bursche,<sup>40</sup> G. Busetto,<sup>22,e</sup> J. Buytaert,<sup>38</sup> S. Cadeddu,<sup>15</sup> R. Calabrese,<sup>16,b</sup> M. Calvi,<sup>20,f</sup> M. Calvo Gomez,<sup>36,g</sup> P. Campana,<sup>18,38</sup> D. Campora Perez,<sup>38</sup> A. Carbone,<sup>14,h</sup> G. Carboni,<sup>24,i</sup> R. Cardinale,<sup>19,38,j</sup> A. Cardini,<sup>15</sup> L. Carson,<sup>50</sup> K. Carvalho Akiba,<sup>2</sup> G. Casse,<sup>52</sup> L. Cassina,<sup>20</sup> L. Castillo Garcia,<sup>38</sup> M. Cattaneo,<sup>38</sup> C. Cauet,<sup>9</sup> R. Cenci,<sup>58</sup> M. Charles,<sup>8</sup> P. Charpentier,<sup>38</sup> S. Chen,<sup>54</sup> S.-F. Cheung,<sup>55</sup> N. Chiapolini,<sup>40</sup> M. Chrzasczcz,<sup>40,26</sup> K. Ciba,<sup>38</sup> X. Cid Vidal,<sup>38</sup> G. Ciezarek,<sup>53</sup> P. E. L. Clarke,<sup>50</sup> M. Clemencic,<sup>38</sup> H. V. Cliff,<sup>47</sup> J. Closier,<sup>38</sup> V. Coco,<sup>38</sup> J. Cogan,<sup>6</sup> E. Cogneras,<sup>5</sup> P. Collins,<sup>38</sup> A. Comerma-Montells,<sup>11</sup> A. Contu,<sup>15</sup> A. Cook,<sup>46</sup> M. Coombes,<sup>46</sup> S. Coquereau,<sup>8</sup> G. Corti,<sup>38</sup> M. Corvo,<sup>16,b</sup> I. Counts,<sup>56</sup> B. Couturier,<sup>38</sup> G. A. Cowan,<sup>50</sup> D. C. Craik,<sup>48</sup> M. Cruz Torres,<sup>60</sup> S. Cunliffe,<sup>53</sup> R. Currie,<sup>50</sup> C. D'Ambrosio,<sup>38</sup> J. Dalseno,<sup>46</sup> P. David,<sup>8</sup> P. N. Y. David,<sup>41</sup> A. Davis,<sup>57</sup> K. De Bruyn,<sup>41</sup> S. De Capua,<sup>54</sup> M. De Cian,<sup>11</sup> J. M. De Miranda,<sup>1</sup> L. De Paula,<sup>2</sup> W. De Silva,<sup>57</sup> P. De Simone,<sup>18</sup> D. Decamp,<sup>4</sup> M. Deckenhoff,<sup>9</sup> L. Del Buono,<sup>8</sup> N. Déleage,<sup>4</sup> D. Derkach,<sup>55</sup> O. Deschamps,<sup>5</sup> F. Dettori,<sup>38</sup> A. Di Canto,<sup>38</sup> H. Dijkstra,<sup>38</sup> S. Donleavy,<sup>52</sup> F. Dordei,<sup>11</sup> M. Dorigo,<sup>39</sup> A. Dosil Suárez,<sup>37</sup> D. Dossett,<sup>48</sup> A. Dovbnya,<sup>43</sup> K. Dreimanis,<sup>52</sup> G. Dujany,<sup>54</sup> F. Dupertuis,<sup>39</sup> P. Durante,<sup>38</sup> R. Dzhelyadin,<sup>35</sup> A. Dziurda,<sup>26</sup> A. Dzyuba,<sup>30</sup> S. Easo,<sup>49,38</sup> U. Egede,<sup>53</sup> V. Egorychev,<sup>31</sup> S. Eidelman,<sup>34</sup> S. Eisenhardt,<sup>50</sup> U. Eitschberger,<sup>9</sup> R. Ekelhof,<sup>9</sup> L. Eklund,<sup>51,38</sup> I. El Rifai,<sup>5</sup> C. Elsasser,<sup>40</sup> S. Ely,<sup>59</sup> S. Esen,<sup>11</sup> H.-M. Evans,<sup>47</sup> T. Evans,<sup>55</sup> A. Falabella,<sup>14</sup> C. Färber,<sup>11</sup> C. Farinelli,<sup>41</sup> N. Farley,<sup>45</sup> S. Farry,<sup>52</sup> R. F. Fay,<sup>52</sup> D. Ferguson,<sup>50</sup> V. Fernandez Albor,<sup>37</sup> F. Ferreira Rodrigues,<sup>1</sup> M. Ferro-Luzzi,<sup>38</sup> S. Filippov,<sup>33</sup> M. Fiore,<sup>16,b</sup> M. Fiorini,<sup>16,b</sup> M. Firlej,<sup>27</sup> C. Fitzpatrick,<sup>38</sup> T. Fiutowski,<sup>27</sup> M. Fontana,<sup>10</sup> F. Fontanelli,<sup>19,j</sup> R. Forty,<sup>38</sup> O. Francisco,<sup>2</sup> M. Frank,<sup>38</sup> C. Frei,<sup>38</sup> M. Frosini,<sup>17,38,a</sup> J. Fu,<sup>21,38</sup> E. Furfaro,<sup>24,i</sup> A. Gallas Torreira,<sup>37</sup> D. Galli,<sup>14,h</sup> S. Gallorini,<sup>22</sup> S. Gambetta,<sup>19,j</sup> M. Gandelman,<sup>2</sup> P. Gandini,<sup>59</sup> Y. Gao,<sup>3</sup> J. García Pardiñas,<sup>37</sup> J. Garofoli,<sup>59</sup> J. Garra Tico,<sup>47</sup> L. Garrido,<sup>36</sup> C. Gaspar,<sup>38</sup> R. Gauld,<sup>55</sup> L. Gavardi,<sup>9</sup> G. Gavrilo, <sup>30</sup> E. Gersabeck,<sup>11</sup> M. Gersabeck,<sup>54</sup> T. Gershon,<sup>48</sup> P. Ghez,<sup>4</sup> A. Gianelle,<sup>22</sup> S. Giani,<sup>39</sup> V. Gibson,<sup>47</sup>

- L. Giubega,<sup>29</sup> V. V. Gligorov,<sup>38</sup> C. Göbel,<sup>60</sup> D. Golubkov,<sup>31</sup> A. Golutvin,<sup>53,31,38</sup> A. Gomes,<sup>1,k</sup> H. Gordon,<sup>38</sup> C. Gotti,<sup>20</sup> M. Grabalosa Gándara,<sup>5</sup> R. Graciani Diaz,<sup>36</sup> L. A. Granado Cardoso,<sup>38</sup> E. Graugés,<sup>36</sup> G. Graziani,<sup>17</sup> A. Grecu,<sup>29</sup> E. Greening,<sup>55</sup> S. Gregson,<sup>47</sup> P. Griffith,<sup>45</sup> L. Grillo,<sup>11</sup> O. Grünberg,<sup>62</sup> B. Gui,<sup>59</sup> E. Gushchin,<sup>33</sup> Y. Guz,<sup>35,38</sup> T. Gys,<sup>38</sup> C. Hadjivasiliou,<sup>59</sup> G. Haefeli,<sup>39</sup> C. Haen,<sup>38</sup> S. C. Haines,<sup>47</sup> S. Hall,<sup>53</sup> B. Hamilton,<sup>58</sup> T. Hampson,<sup>46</sup> X. Han,<sup>11</sup> S. Hansmann-Menzemer,<sup>11</sup> N. Harnew,<sup>55</sup> S. T. Harnew,<sup>46</sup> J. Harrison,<sup>54</sup> J. He,<sup>38</sup> T. Head,<sup>38</sup> V. Heijne,<sup>41</sup> K. Hennessy,<sup>52</sup> P. Henrard,<sup>5</sup> L. Henry,<sup>8</sup> J. A. Hernando Morata,<sup>37</sup> E. van Herwijnen,<sup>38</sup> M. Heß,<sup>62</sup> A. Hicheur,<sup>1</sup> D. Hill,<sup>55</sup> M. Hoballah,<sup>5</sup> C. Hombach,<sup>54</sup> W. Hulsbergen,<sup>41</sup> P. Hunt,<sup>55</sup> N. Hussain,<sup>55</sup> D. Hutchcroft,<sup>52</sup> D. Hynds,<sup>51</sup> M. Idzik,<sup>27</sup> P. Ilten,<sup>56</sup> R. Jacobsson,<sup>38</sup> A. Jaeger,<sup>11</sup> J. Jalocha,<sup>55</sup> E. Jans,<sup>41</sup> P. Jaton,<sup>39</sup> A. Jawahery,<sup>58</sup> F. Jing,<sup>3</sup> M. John,<sup>55</sup> D. Johnson,<sup>55</sup> C. R. Jones,<sup>47</sup> C. Joram,<sup>38</sup> B. Jost,<sup>38</sup> N. Jurik,<sup>59</sup> M. Kabbalo,<sup>9</sup> S. Kandybei,<sup>43</sup> W. Kanso,<sup>6</sup> M. Karacson,<sup>38</sup> T. M. Karbach,<sup>38</sup> S. Karodia,<sup>51</sup> M. Kelsey,<sup>59</sup> I. R. Kenyon,<sup>45</sup> T. Ketel,<sup>42</sup> B. Khanji,<sup>20</sup> C. Khurewathanakul,<sup>39</sup> S. Klaver,<sup>54</sup> K. Klimaszewski,<sup>28</sup> O. Kochebina,<sup>7</sup> M. Kolpin,<sup>11</sup> I. Komarov,<sup>39</sup> R. F. Koopman,<sup>42</sup> P. Koppenburg,<sup>41,38</sup> M. Korolev,<sup>32</sup> A. Kozlinskiy,<sup>41</sup> L. Kravchuk,<sup>33</sup> K. Kreplin,<sup>11</sup> M. Kreps,<sup>48</sup> G. Krocker,<sup>11</sup> P. Krokovny,<sup>34</sup> F. Kruse,<sup>9</sup> W. Kucewicz,<sup>26,l</sup> M. Kucharczyk,<sup>20,26,38,f</sup> V. Kudryavtsev,<sup>34</sup> K. Kurek,<sup>28</sup> T. Kvaratskheliya,<sup>31</sup> V. N. La Thi,<sup>39</sup> D. Lacarrere,<sup>38</sup> G. Lafferty,<sup>54</sup> A. Lai,<sup>15</sup> D. Lambert,<sup>50</sup> R. W. Lambert,<sup>42</sup> G. Lanfranchi,<sup>18</sup> C. Langenbruch,<sup>48</sup> B. Langhans,<sup>38</sup> T. Latham,<sup>48</sup> C. Lazzeroni,<sup>45</sup> R. Le Gac,<sup>6</sup> J. van Leerdam,<sup>41</sup> J.-P. Lees,<sup>4</sup> R. Lefèvre,<sup>5</sup> A. Leflat,<sup>32</sup> J. Lefrançois,<sup>7</sup> S. Leo,<sup>23</sup> O. Leroy,<sup>6</sup> T. Lesiak,<sup>26</sup> B. Leverington,<sup>11</sup> Y. Li,<sup>3</sup> T. Likhomanenko,<sup>63</sup> M. Liles,<sup>52</sup> R. Lindner,<sup>38</sup> C. Linn,<sup>38</sup> F. Lionetto,<sup>40</sup> B. Liu,<sup>15</sup> G. Liu,<sup>38</sup> S. Lohn,<sup>38</sup> I. Longstaff,<sup>51</sup> J. H. Lopes,<sup>2</sup> N. Lopez-March,<sup>39</sup> P. Lowdon,<sup>40</sup> H. Lu,<sup>3</sup> D. Lucchesi,<sup>22,e</sup> H. Luo,<sup>50</sup> A. Lupato,<sup>22</sup> E. Luppi,<sup>16,b</sup> O. Lupton,<sup>55</sup> F. Machefert,<sup>7</sup> I. V. Machikhiliyan,<sup>31</sup> F. Maciuc,<sup>29</sup> O. Maev,<sup>30</sup> S. Malde,<sup>55</sup> G. Manca,<sup>15,m</sup> G. Mancinelli,<sup>6</sup> J. Maratas,<sup>5</sup> J. F. Marchand,<sup>4</sup> U. Marconi,<sup>14</sup> C. Marin Benito,<sup>36</sup> P. Marino,<sup>23,n</sup> R. Märki,<sup>39</sup> J. Marks,<sup>11</sup> G. Martellotti,<sup>25</sup> A. Martens,<sup>8</sup> A. Martín Sánchez,<sup>7</sup> M. Martinelli,<sup>41</sup> D. Martinez Santos,<sup>42</sup> F. Martinez Vidal,<sup>64</sup> D. Martins Tostes,<sup>2</sup> A. Massafferri,<sup>1</sup> R. Matev,<sup>38</sup> Z. Mathe,<sup>38</sup> C. Matteuzzi,<sup>20</sup> A. Mazurov,<sup>16,b</sup> M. McCann,<sup>53</sup> J. McCarthy,<sup>45</sup> A. McNab,<sup>54</sup> R. McNulty,<sup>12</sup> B. McKelley,<sup>52</sup> B. Meadows,<sup>57</sup> F. Meier,<sup>9</sup> M. Meissner,<sup>11</sup> M. Merk,<sup>41</sup> D. A. Milanes,<sup>8</sup> M.-N. Minard,<sup>4</sup> N. Moggi,<sup>14</sup> J. Molina Rodriguez,<sup>60</sup> S. Monteil,<sup>5</sup> M. Morandin,<sup>22</sup> P. Morawski,<sup>27</sup> A. Mordà,<sup>6</sup> M. J. Morello,<sup>23,n</sup> J. Moron,<sup>27</sup> A.-B. Morris,<sup>50</sup> R. Mountain,<sup>59</sup> F. Muheim,<sup>50</sup> K. Müller,<sup>40</sup> M. Mussini,<sup>14</sup> B. Muster,<sup>39</sup> P. Naik,<sup>46</sup> T. Nakada,<sup>39</sup> R. Nandakumar,<sup>49</sup> I. Nasteva,<sup>2</sup> M. Needham,<sup>50</sup> N. Neri,<sup>21</sup> S. Neubert,<sup>38</sup> N. Neufeld,<sup>38</sup> M. Neuner,<sup>11</sup> A. D. Nguyen,<sup>39</sup> T. D. Nguyen,<sup>39</sup> C. Nguyen-Mau,<sup>39,o</sup> M. Nicol,<sup>7</sup> V. Niess,<sup>5</sup> R. Niet,<sup>9</sup> N. Nikitin,<sup>32</sup> T. Nikodem,<sup>11</sup> A. Novoselov,<sup>35</sup> D. P. O'Hanlon,<sup>48</sup> A. Oblakowska-Mucha,<sup>27</sup> V. Obraztsov,<sup>35</sup> S. Oggero,<sup>41</sup> S. Ogilvy,<sup>51</sup> O. Okhrimenko,<sup>44</sup> R. Oldeman,<sup>15,m</sup> G. Onderwater,<sup>65</sup> M. Orlandea,<sup>29</sup> J. M. Otalora Goicochea,<sup>2</sup> P. Owen,<sup>53</sup> A. Oyanguren,<sup>64</sup> B. K. Pal,<sup>59</sup> A. Palano,<sup>13,p</sup> F. Palombo,<sup>21,q</sup> M. Palutan,<sup>18</sup> J. Panman,<sup>38</sup> A. Papanestis,<sup>49,38</sup> M. Pappagallo,<sup>51</sup> C. Parkes,<sup>54</sup> C. J. Parkinson,<sup>9,45</sup> G. Passaleva,<sup>17</sup> G. D. Patel,<sup>52</sup> M. Patel,<sup>53</sup> C. Patrignani,<sup>19,j</sup> A. Pazos Alvarez,<sup>37</sup> A. Pearce,<sup>54</sup> A. Pellegrino,<sup>41</sup> M. Pepe Altarelli,<sup>38</sup> S. Perazzini,<sup>14,h</sup> E. Perez Trigo,<sup>37</sup> P. Perret,<sup>5</sup> M. Perrin-Terrin,<sup>6</sup> L. Pescatore,<sup>45</sup> E. Pesen,<sup>66</sup> K. Petridis,<sup>53</sup> A. Petrolini,<sup>19,j</sup> E. Picatoste Olloqui,<sup>36</sup> B. Pietrzyk,<sup>4</sup> T. Pilař,<sup>48</sup> D. Pinci,<sup>25</sup> A. Pistone,<sup>19</sup> S. Playfer,<sup>50</sup> M. Plo Casasus,<sup>37</sup> F. Polci,<sup>8</sup> A. Poluektov,<sup>48,34</sup> E. Polycarpo,<sup>2</sup> A. Popov,<sup>35</sup> D. Popov,<sup>10</sup> B. Popovici,<sup>29</sup> C. Potterat,<sup>2</sup> E. Price,<sup>46</sup> J. Prisciandaro,<sup>39</sup> A. Pritchard,<sup>52</sup> C. Prouve,<sup>46</sup> V. Pugatch,<sup>44</sup> A. Puig Navarro,<sup>39</sup> G. Punzi,<sup>23,r</sup> W. Qian,<sup>4</sup> B. Rachwal,<sup>26</sup> J. H. Rademacker,<sup>46</sup> B. Rakotomiramanana,<sup>39</sup> M. Rama,<sup>18</sup> M. S. Rangel,<sup>2</sup> I. Raniuk,<sup>43</sup> N. Rauschmayr,<sup>38</sup> G. Raven,<sup>42</sup> S. Reichert,<sup>54</sup> M. M. Reid,<sup>48</sup> A. C. dos Reis,<sup>1</sup> S. Ricciardi,<sup>49</sup> S. Richards,<sup>46</sup> M. Rihl,<sup>38</sup> K. Rinnert,<sup>52</sup> V. Rives Molina,<sup>36</sup> D. A. Roa Romero,<sup>5</sup> P. Robbe,<sup>7</sup> A. B. Rodrigues,<sup>1</sup> E. Rodrigues,<sup>54</sup> P. Rodriguez Perez,<sup>54</sup> S. Roiser,<sup>38</sup> V. Romanovsky,<sup>35</sup> A. Romero Vidal,<sup>37</sup> M. Rotondo,<sup>22</sup> J. Rouvinet,<sup>39</sup> T. Ruf,<sup>38</sup> F. Ruffini,<sup>23</sup> H. Ruiz,<sup>36</sup> P. Ruiz Valls,<sup>64</sup> J. J. Saborido Silva,<sup>37</sup> N. Sagidova,<sup>30</sup> P. Sail,<sup>51</sup> B. Saitta,<sup>15,m</sup> V. Salustino Guimaraes,<sup>2</sup> C. Sanchez Mayordomo,<sup>64</sup> B. Sanmartin Sedes,<sup>37</sup> R. Santacesaria,<sup>25</sup> C. Santamarina Rios,<sup>37</sup> E. Santovetti,<sup>24,i</sup> A. Sarti,<sup>18,s</sup> C. Satriano,<sup>25,c</sup> A. Satta,<sup>24</sup> D. M. Saunders,<sup>46</sup> M. Savrie,<sup>16,b</sup> D. Savrina,<sup>31,32</sup> M. Schiller,<sup>42</sup> H. Schindler,<sup>38</sup> M. Schlupp,<sup>9</sup> M. Schmelling,<sup>10</sup> B. Schmidt,<sup>38</sup> O. Schneider,<sup>39</sup> A. Schopper,<sup>38</sup> M.-H. Schune,<sup>7</sup> R. Schwemmer,<sup>38</sup> B. Sciascia,<sup>18</sup> A. Sciubba,<sup>25</sup> M. Seco,<sup>37</sup> A. Semennikov,<sup>31</sup> I. Sepp,<sup>53</sup> N. Serra,<sup>40</sup> J. Serrano,<sup>6</sup> L. Sestini,<sup>22</sup> P. Seyfert,<sup>11</sup> M. Shapkin,<sup>35</sup> I. Shapoval,<sup>16,43,b</sup> Y. Shcheglov,<sup>30</sup> T. Shears,<sup>52</sup> L. Shekhtman,<sup>34</sup> V. Shevchenko,<sup>63</sup> A. Shires,<sup>9</sup> R. Silva Coutinho,<sup>48</sup> G. Simi,<sup>22</sup> M. Sirendi,<sup>47</sup> N. Skidmore,<sup>46</sup> T. Skwarnicki,<sup>59</sup> N. A. Smith,<sup>52</sup> E. Smith,<sup>55,49</sup> E. Smith,<sup>53</sup> J. Smith,<sup>47</sup> M. Smith,<sup>54</sup> H. Snoek,<sup>41</sup> M. D. Sokoloff,<sup>57</sup> F. J. P. Soler,<sup>51</sup> F. Soomro,<sup>39</sup> D. Souza,<sup>46</sup> B. Souza De Paula,<sup>2</sup> B. Spaan,<sup>9</sup> A. Sparkes,<sup>50</sup> P. Spradlin,<sup>51</sup> S. Sridharan,<sup>38</sup> F. Stagni,<sup>38</sup> M. Stahl,<sup>11</sup> S. Stahl,<sup>11</sup> O. Steinkamp,<sup>40</sup> O. Stenyakin,<sup>35</sup> S. Stevenson,<sup>55</sup> S. Stoica,<sup>29</sup> S. Stone,<sup>59</sup> B. Storaci,<sup>40</sup> S. Stracka,<sup>23,38</sup> M. Straticiu,<sup>29</sup> U. Straumann,<sup>40</sup> R. Stroili,<sup>22</sup> V. K. Subbiah,<sup>38</sup> L. Sun,<sup>57</sup> W. Sutcliffe,<sup>53</sup> K. Swientek,<sup>27</sup> S. Swientek,<sup>9</sup> V. Syropoulos,<sup>42</sup> M. Szczekowski,<sup>28</sup> P. Szczypka,<sup>39,38</sup> D. Szilard,<sup>2</sup> T. Szumlak,<sup>27</sup> S. T'Jampens,<sup>4</sup> M. Teklishyn,<sup>7</sup> G. Tellarini,<sup>16,b</sup> F. Teubert,<sup>38</sup>

C. Thomas,<sup>55</sup> E. Thomas,<sup>38</sup> J. van Tilburg,<sup>41</sup> V. Tisserand,<sup>4</sup> M. Tobin,<sup>39</sup> S. Tolk,<sup>42</sup> L. Tomassetti,<sup>16,b</sup> D. Tonelli,<sup>38</sup> S. Topp-Joergensen,<sup>55</sup> N. Torr,<sup>55</sup> E. Tournefier,<sup>4</sup> S. Tourneur,<sup>39</sup> M. T. Tran,<sup>39</sup> M. Tresch,<sup>40</sup> A. Tsaregorodtsev,<sup>6</sup> P. Tsopelas,<sup>41</sup> N. Tuning,<sup>41</sup> M. Ubeda Garcia,<sup>38</sup> A. Ukleja,<sup>28</sup> A. Ustyuzhanin,<sup>63</sup> U. Uwer,<sup>11</sup> V. Vagnoni,<sup>14</sup> G. Valenti,<sup>14</sup> A. Vallier,<sup>7</sup> R. Vazquez Gomez,<sup>18</sup> P. Vazquez Regueiro,<sup>37</sup> C. Vázquez Sierra,<sup>37</sup> S. Vecchi,<sup>16</sup> J. J. Velthuis,<sup>46</sup> M. Veltri,<sup>17,t</sup> G. Veneziano,<sup>39</sup> M. Vesterinen,<sup>11</sup> B. Viaud,<sup>7</sup> D. Vieira,<sup>2</sup> M. Vieites Diaz,<sup>37</sup> X. Vilasis-Cardona,<sup>36,g</sup> A. Vollhardt,<sup>40</sup> D. Volyanskyy,<sup>10</sup> D. Voong,<sup>46</sup> A. Vorobyev,<sup>30</sup> V. Vorobyev,<sup>34</sup> C. Voß,<sup>62</sup> H. Voss,<sup>10</sup> J. A. de Vries,<sup>41</sup> R. Waldi,<sup>62</sup> C. Wallace,<sup>48</sup> R. Wallace,<sup>12</sup> J. Walsh,<sup>23</sup> S. Wandernoth,<sup>11</sup> J. Wang,<sup>59</sup> D. R. Ward,<sup>47</sup> N. K. Watson,<sup>45</sup> D. Websdale,<sup>53</sup> M. Whitehead,<sup>48</sup> J. Wicht,<sup>38</sup> D. Wiedner,<sup>11</sup> G. Wilkinson,<sup>55</sup> M. P. Williams,<sup>45</sup> M. Williams,<sup>56</sup> F. F. Wilson,<sup>49</sup> J. Wimberley,<sup>58</sup> J. Wishahi,<sup>9</sup> W. Wislicki,<sup>28</sup> M. Witek,<sup>26</sup> G. Wormser,<sup>7</sup> S. A. Wotton,<sup>47</sup> S. Wright,<sup>47</sup> S. Wu,<sup>3</sup> K. Wyllie,<sup>38</sup> Y. Xie,<sup>61</sup> Z. Xing,<sup>59</sup> Z. Xu,<sup>39</sup> Z. Yang,<sup>3</sup> X. Yuan,<sup>3</sup> O. Yushchenko,<sup>35</sup> M. Zangoli,<sup>14</sup> M. Zavertyaev,<sup>10,u</sup> L. Zhang,<sup>59</sup> W. C. Zhang,<sup>12</sup> Y. Zhang,<sup>3</sup> A. Zhelezov,<sup>11</sup> A. Zhokhov,<sup>31</sup> L. Zhong,<sup>3</sup> and A. Zvyagin<sup>38</sup>

## (LHCb Collaboration)

- <sup>1</sup>*Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil*  
<sup>2</sup>*Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil*  
<sup>3</sup>*Center for High Energy Physics, Tsinghua University, Beijing, China*  
<sup>4</sup>*LAPP, Université de Savoie, CNRS/IN2P3, Annecy-Le-Vieux, France*  
<sup>5</sup>*Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France*  
<sup>6</sup>*CPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France*  
<sup>7</sup>*LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France*  
<sup>8</sup>*LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France*  
<sup>9</sup>*Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*  
<sup>10</sup>*Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*  
<sup>11</sup>*Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*  
<sup>12</sup>*School of Physics, University College Dublin, Dublin, Ireland*  
<sup>13</sup>*Sezione INFN di Bari, Bari, Italy*  
<sup>14</sup>*Sezione INFN di Bologna, Bologna, Italy*  
<sup>15</sup>*Sezione INFN di Cagliari, Cagliari, Italy*  
<sup>16</sup>*Sezione INFN di Ferrara, Ferrara, Italy*  
<sup>17</sup>*Sezione INFN di Firenze, Firenze, Italy*  
<sup>18</sup>*Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy*  
<sup>19</sup>*Sezione INFN di Genova, Genova, Italy*  
<sup>20</sup>*Sezione INFN di Milano Bicocca, Milano, Italy*  
<sup>21</sup>*Sezione INFN di Milano, Milano, Italy*  
<sup>22</sup>*Sezione INFN di Padova, Padova, Italy*  
<sup>23</sup>*Sezione INFN di Pisa, Pisa, Italy*  
<sup>24</sup>*Sezione INFN di Roma Tor Vergata, Roma, Italy*  
<sup>25</sup>*Sezione INFN di Roma La Sapienza, Roma, Italy*  
<sup>26</sup>*Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*  
<sup>27</sup>*AGH–University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*  
<sup>28</sup>*National Center for Nuclear Research (NCBJ), Warsaw, Poland*  
<sup>29</sup>*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*  
<sup>30</sup>*Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*  
<sup>31</sup>*Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*  
<sup>32</sup>*Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*  
<sup>33</sup>*Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*  
<sup>34</sup>*Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*  
<sup>35</sup>*Institute for High Energy Physics (IHEP), Protvino, Russia*  
<sup>36</sup>*Universitat de Barcelona, Barcelona, Spain*  
<sup>37</sup>*Universidad de Santiago de Compostela, Santiago de Compostela, Spain*  
<sup>38</sup>*European Organization for Nuclear Research (CERN), Geneva, Switzerland*  
<sup>39</sup>*Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*  
<sup>40</sup>*Physik-Institut, Universität Zürich, Zürich, Switzerland*  
<sup>41</sup>*Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*

- <sup>42</sup>*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- <sup>43</sup>*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>44</sup>*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>45</sup>*University of Birmingham, Birmingham, United Kingdom*
- <sup>46</sup>*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>47</sup>*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>48</sup>*Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>49</sup>*STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>50</sup>*School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>51</sup>*School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>52</sup>*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>53</sup>*Imperial College London, London, United Kingdom*
- <sup>54</sup>*School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>55</sup>*Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>56</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts, United States*
- <sup>57</sup>*University of Cincinnati, Cincinnati, Ohio, United States*
- <sup>58</sup>*University of Maryland, College Park, Maryland, United States*
- <sup>59</sup>*Syracuse University, Syracuse, New York, United States*
- <sup>60</sup>*Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil (associated with Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil)*
- <sup>61</sup>*Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China (associated with Center for High Energy Physics, Tsinghua University, Beijing, China)*
- <sup>62</sup>*Institut für Physik, Universität Rostock, Rostock, Germany (associated with Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany)*
- <sup>63</sup>*National Research Centre Kurchatov Institute, Moscow, Russia (associated with Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia)*
- <sup>64</sup>*Instituto de Física Corpuscular (IFIC), Universitat de Valencia-CSIC, Valencia, Spain (associated with Universitat de Barcelona, Barcelona, Spain)*
- <sup>65</sup>*KVI–University of Groningen, Groningen, The Netherlands (associated with Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands)*
- <sup>66</sup>*Celal Bayar University, Manisa, Turkey (associated with European Organization for Nuclear Research (CERN), Geneva, Switzerland)*

<sup>a</sup>Also at Università di Firenze, Firenze, Italy.

<sup>b</sup>Also at Università di Ferrara, Ferrara, Italy.

<sup>c</sup>Also at Università della Basilicata, Potenza, Italy.

<sup>d</sup>Also at Università di Modena e Reggio Emilia, Modena, Italy.

<sup>e</sup>Also at Università di Padova, Padova, Italy.

<sup>f</sup>Also at Università di Milano Bicocca, Milano, Italy.

<sup>g</sup>Also at LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain.

<sup>h</sup>Also at Università di Bologna, Bologna, Italy.

<sup>i</sup>Also at Università di Roma Tor Vergata, Roma, Italy.

<sup>j</sup>Also at Università di Genova, Genova, Italy.

<sup>k</sup>Also at Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil.

<sup>l</sup>Also at AGH–University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland.

<sup>m</sup>Also at Università di Cagliari, Cagliari, Italy.

<sup>n</sup>Also at Scuola Normale Superiore, Pisa, Italy.

<sup>o</sup>Also at Hanoi University of Science, Hanoi, Viet Nam.

<sup>p</sup>Also at Università di Bari, Bari, Italy.

<sup>q</sup>Also at Università degli Studi di Milano, Milano, Italy.

<sup>r</sup>Also at Università di Pisa, Pisa, Italy.

<sup>s</sup>Also at Università di Roma La Sapienza, Roma, Italy.

<sup>t</sup>Also at Università di Urbino, Urbino, Italy.

<sup>u</sup>Also at P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia.